

INVESTOR IN PEOPLE

**BEST AVAILABLE COPY**

# **PRIORITY DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)


The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 15 July 2004

THE PATENT OFFICE  
A

14 AUG 2003

LONDON

15AUG03 EB30509-1 001030  
P01/7700 0.00-0319150.9

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office  
Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference

BJR/JW/PB60445

2. Patent application number

(The Patent Office will fill in his part)

0319150.9

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

Glaxo Group Limited  
Glaxo Wellcome House, Berkeley Avenue,  
Greenford, Middlesex UB6 0NN, Great Britain

473587003  
United Kingdom

4. Title of the invention

Novel Compounds

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Patents ADP number (if you know it)

Corporate Intellectual Property

GlaxoSmithKline  
Corporate Intellectual Property (CN9 25.1)  
980 Great West Road  
BRENTFORD  
Middlesex TW8 9GS

8672555006

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number

Country	Priority application number (if you know it)	Date of filing (day / month / year)
---------	--	-------------------------------------

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application	Date of filing (day / month / year)
-------------------------------	-------------------------------------

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer yes if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is named as an applicant, or
  - c) any named applicant is a corporate body
- See note (d)

**Patents Form 1/77**

Enter the number of sheets for any of the following items you are filing with this form.  
Do not count copies of the same document

Continuation sheets of this form  
Description  
Claim(s)  
Abstract  
Drawings

20 ✓  
2 ✓

CF

10. If you are also filing any of the following, state how many against each item.

**Priority Documents**

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents  
(please specify)

11.

We request the grant of a patent on the basis of this application

Signature

B J Russell

Date 14-Aug-03

12. Name and daytime telephone number of person to contact in the United Kingdom

B J Russell 01279 644398

**Warning**

*After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- f) For details of the fee and ways to pay please contact the Patent Office.

## Novel Compounds

5 This invention relates to novel amide derivatives having pharmacological activity, processes for their preparation, to compositions containing them and to their use in medicine, especially in the treatment of various disorders.

Vanilloids are a class of natural and synthetic compounds that are characterised by the presence of a vanillyl (4-hydroxy 3-methoxybenzyl) group or a  
10 functionally equivalent group. Vanilloid Receptor (VR-1), whose function is modulated by such compounds, has been widely studied and is extensively reviewed by Szallasi and Blumberg (The American Society for Pharmacology and Experimental Therapeutics, 1999, Vol. 51, No. 2.).

A wide variety of Vanilloid compounds of different structures are known  
15 in the art, for example those disclosed in European Patent Application Numbers, EP 0 347 000 and EP 0 401 903, UK Patent Application Number GB 2226313 and International Patent Applications, Publication Numbers WO 92/09285, WO 02/100819 and WO 02/076946. Particularly notable examples of vanilloid compounds or vanilloid receptor modulators are capsaicin or trans 8-methyl-N-vanillyl-6-nonenamide which is isolated from the pepper plant, capsazepine  
20 (*Tetrahedron*, 53, 1997, 4791) and olvanil or - N-(4-hydroxy-3-methoxybenzyl)oleamide (*J. Med. Chem.*, 36, 1993, 2595).


International Patent Application, Publication Number WO 02/08221 discloses diaryl piperazine and related compounds which bind with high  
25 selectivity and high affinity to vanilloid receptors, especially Type I Vanilloid receptors, also known as capsaicin or VR1 receptors. The compounds are said to be useful in the treatment of chronic and acute pain conditions, itch and urinary incontinence.

International Patent Applications, Publication Numbers WO 02/16317,  
30 WO 02/16318 and WO 02/16319 suggest that compounds having a high affinity for the vanilloid receptor are useful for treating stomach-duodenal ulcers.

International Patent Applications, Publication Numbers WO 02/072536, WO 02/090326, WO 03/022809 and WO 03/053945; and International Patent

Application Number PCT/GB03/00608 also describe a variety of compounds having activity as vanilloid receptor antagonists.

According to a first aspect of the present invention, there is provided a compound of formula (I),



or a pharmaceutically acceptable salt or solvate thereof, wherein,

10 P represents phenyl, quinoliny, isoquinoliny, 1,2,3,4-tetrahydroquinoliny, benzoisoxazolyl or benzothiazolyl;

P' represents phenyl, pyridinyl or pyrimidinyl;

R¹ and R³ may be the same or different and represent alkyl, alkoxy, halo, -CF₃, -OCF₃, -OH, =O, -CN, -NO₂, -SO₂NH₂, -SO₂R⁴ or -NR⁴R⁵;

15 R² represents -H;

R⁴ and R⁵ may be the same or different and represent -H or alkyl;

m represents 0 or 1;

n represents 0, 1, 2, 3, 4 or 5; and

X represents N or CH.

20

Preferably, P represents phenyl. Preferably, P represents quinoliny, isoquinoliny, 1,2,3,4-tetrahydroquinoliny, benzoisoxazolyl or benzothiazolyl.

Preferably, P' represents phenyl. Preferably, P' represents pyridinyl or pyrimidinyl.

25

Preferably, R¹ represents alkyl, halo, =O, -SO₂NH₂, or -SO₂Me.

Preferably, R³ represents alkyl, alkoxy, halo, -CF₃ or -CN.

Preferably, R⁴ is -H or methyl.



Preferably, R<sup>5</sup> is -H or methyl.

Preferably, m represents 0. Preferably, m represents 1.

Preferably, n represents 0, 1 or 2.

Preferably, X represents N. Preferably, X represents CH.

5 Preferred compounds according to this invention include Examples 1 - 40 or pharmaceutically acceptable salts or solvates thereof.

Certain of the carbon atoms of formula (I) are chiral carbon atoms, and therefore compounds of formula (I) may exist as stereoisomers. The invention extends to all optical isomers such as stereoisomeric forms of the compounds of  
10 formula (I) including enantiomers and mixtures thereof, such as racemates. The different stereoisomeric forms may be separated or resolved one from the other by conventional methods or any given isomer may be obtained by conventional stereospecific or asymmetric syntheses.

As indicated above, the compounds of formula (I) can form salts,  
15 especially pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts are those used conventionally in the art and include those described in *J. Pharm. Sci.*, 1977, 66, 1-19, such as acid addition salts.

Suitable pharmaceutically acceptable salts include acid addition salts.

Suitable pharmaceutically acceptable acid addition salts include salts with  
20 inorganic acids such, for example, as hydrochloric acid, hydrobromic acid, orthophosphoric acid or sulphuric acid, or with organic acids such, for example as methanesulphonic acid, toluenesulphonic acid, acetic acid, propionic acid, lactic acid, citric acid, fumaric acid, malic acid, succinic acid, salicylic acid, maleic acid, glycerophosphoric acid or acetylsalicylic acid.

25 The salts and/or solvates of the compounds of the formula (I) which are not pharmaceutically acceptable may be useful as intermediates in the preparation of pharmaceutically acceptable salts and/or solvates of compounds of formula (I) or the compounds of the formula (I) themselves, and as such form another aspect of the present invention.

30 The compounds of formula (I) may be prepared in crystalline or non-crystalline form, and if crystalline, may be optionally hydrated or solvated. This invention includes in its scope stoichiometric hydrates as well as compounds containing variable amounts of water.

Suitable solvates include pharmaceutically acceptable solvates, such as hydrates.

Solvates include stoichiometric solvates and non-stoichiometric solvates.


As used herein the term "alkyl" as a group or part of a group refers to a  
5 straight or branched chain saturated aliphatic hydrocarbon radical containing 1 to  
12 carbon atoms, suitably 1 to 6 carbon atoms. Such alkyl groups in particular  
include methyl ("Me"), ethyl ("Et"), n-propyl ("Pr<sup>n</sup>"), *iso*-propyl ("Pr<sup>i</sup>"), n-butyl  
("Bu<sup>n</sup>"), *sec*-butyl ("Bu<sup>s</sup>"), *tert*-butyl ("Bu<sup>t</sup>"), pentyl and hexyl. The term  
"cycloalkyl" as part of a group refers to a saturated alicyclic hydrocarbon radical  
10 containing 3 to 12 carbon atoms, suitably 3 to 6 carbon atoms. Where  
appropriate, such alkyl groups may be substituted by one or more groups  
selected from halo (such as fluoro, chloro, bromo), -CN, -CF<sub>3</sub>, -OH, -OCF<sub>3</sub>, C<sub>2</sub>-  
6 alkenyl, C<sub>3</sub>-6 alkynyl, C<sub>1</sub>-6 alkoxy, aryl and di-C<sub>1</sub>-6 alkylamino. Alkyl is  
preferably unsubstituted.

15 As used herein, the term "alkoxy" as a group or part of a group refers to  
an alkyl ether radical, wherein the term "alkyl" is defined above. Such alkoxy  
groups in particular include methoxy, ethoxy, n-propoxy, *iso*-propoxy, n-butoxy,  
*iso*-butoxy, *sec*-butoxy and *tert*-butoxy. Where appropriate, such alkoxy groups  
may be substituted by one or more groups selected from halo (such as fluoro,  
20 chloro, bromo), -CN, -CF<sub>3</sub>, -OH, -OCF<sub>3</sub>, C<sub>1</sub>-6 alkyl, C<sub>2</sub>-6 alkenyl, C<sub>3</sub>-6 alkynyl,  
aryl and di-C<sub>1</sub>-6 alkylamino. Alkoxy is preferably unsubstituted.

The term "halo" is used herein to describe, unless otherwise stated, a  
group selected from fluorine ("fluoro"), chlorine ("chloro"), bromine ("bromo") or  
iodine ("iodo").


25 The present invention also provides a process for the preparation of a  
compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof,  
which process comprises:

(a) reacting a compound of formula (II),



5

wherein, P, R<sup>1</sup> and n are as defined in relation to formula (I), with a compound of formula (III),



10


wherein, P', R<sup>3</sup>, m, n and X are as defined in relation to formula (I) and thereafter, as necessary, carrying out one or more of the following reactions:

- 15 (i) converting one compound of formula (I) into another compound of formula (I);  
(ii) removing any protecting group;  
(iii) preparing a salt or a solvate of the compound so formed.

20 The reaction between a compound of formula (II) and a compound of formula (III) may be effected using conventional methods for the formation of an amide bond, such as those described in J March, *Advanced Organic Chemistry*, 4th edition, J Wiley & Sons, 1992, p. 419-421.

Compounds of formula (II) are either commercially available, or may be prepared by the reaction of a compound of formula (IV),





(IV)


5 wherein, P, R<sup>1</sup> and n are as defined in relation to formula (I), with a suitable reducing agent.

The reaction of a compound of formula (IV) with a reducing agent may be effected by methods well known in the art, such as those described in J March, *Advanced Organic Chemistry*, 4th edition, J Wiley & Sons, 1992, p. 1216-1218.

10 Suitable reducing agents include (a) iron or zinc metal in hydrochloric acid, or (b) hydrogen in the presence of a suitable catalyst, such as, 5% palladium on charcoal. Reduction using hydrogen may conveniently be performed in a solvent such as methanol or ethanol.

15 Compounds of formula (IV) are commercially available or may be prepared according to literature methods.

Compounds of formula (III) are either commercially available or may be prepared by hydrolysis of a compound of formula (V),



(V)

20 wherein, P', R<sup>3</sup>, m, n and X are as defined in relation to formula (I) and R' is an alkyl group. A suitable hydrolysis agent is aqueous hydrochloric acid. A suitable solvent is ethyl acetate.

Compounds of formula (V) are commercially available or may be prepared according to literature methods such as those described in *J. Org. Chem.* 28, 1963, 3259 or *J. Am. Chem. Soc.* 118, 1996, 7215.

The above-mentioned conversions of a compound of formula (I) into another compound of formula (I) include any conversion, which may be effected using conventional procedures, but in particular the said conversions include any combination of:

- (i) converting one group  $R^1$  into another group  $R^1$ ; and
- (ii) converting one group  $R^3$  into another group  $R^3$ .

The above-mentioned conversions (i) – (ii) may be performed using any appropriate method under conditions determined by the particular groups chosen.

It will be appreciated by those skilled in the art that it may be necessary to protect certain reactive substituents during some of the above procedures. Standard protection and deprotection techniques, such as those described in Greene T.W. 'Protective groups in organic synthesis', New York, Wiley (1981), can be used. For example, primary amines can be protected as phthalimide, benzyl, benzyloxycarbonyl or trityl derivatives. Carboxylic acid groups can be protected as esters. Aldehyde or ketone groups can be protected as acetals, ketals, thioacetals or thioketals. Deprotection of such groups is achieved using conventional procedures known in the art.

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

Compounds of formula (I) and their pharmaceutically acceptable salts and solvates thereof have Vanilloid receptor antagonist (VR1) activity and are believed to be of potential use for the treatment or prophylaxis of certain disorders, or treatment of the pain associated with them, such as: pain, chronic pain, neuropathic pain, postoperative pain, post-rheumatoid arthritic pain, osteoarthritic pain, back pain, visceral pain, cancer pain, algesia, neuralgia, dental pain, headache, migraine, neuropathies, carpal tunnel syndrome, diabetic neuropathy, HIV-related neuropathy, post-herpetic neuralgia, fibromyalgia, neuritis, sciatica, nerve injury, ischaemia, neurodegeneration, stroke, post stroke

pain, multiple sclerosis, respiratory diseases, asthma, cough, COPD, broncho constriction, inflammatory disorders, oesophagitis, heart burn, Barrett's metaplasia, dysphagia, gastroesophageal reflux disorder (GERD), stomach and duodenal ulcers, functional dyspepsia, irritable bowel syndrome, inflammatory  
5 bowel disease, colitis, Crohn's disease, pelvic hypersensitivity, pelvic pain, menstrual pain, renal colic, urinary incontinence, cystitis, burns, itch, psoriasis, pruritis, emesis (hereinafter referred to as the "Disorders of the Invention").

Accordingly, the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, for use as an active  
10 therapeutic substance, in particular, in the treatment and/or prophylaxis of the Disorders of the Invention.

In particular, the invention provides a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof for use in the treatment or prophylaxis of pain.

15 The invention further provides a method for the treatment or prophylaxis of disorders in which antagonism of the Vanilloid (VR1) receptor is beneficial, in particular the Disorders of the Invention, in mammals including humans, which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable  
20 salt or solvate thereof.

The invention provides for the use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment or prophylaxis of disorders in which antagonism of the Vanilloid (VR1) receptor is beneficial, particularly the Disorders of the  
25 Invention.

In order to use the compounds of the invention in therapy, they will normally be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice. Thus, the present invention also provides a pharmaceutical composition, which comprises a compound of formula (I) or a  
30 pharmaceutically acceptable salt or solvate thereof and a pharmaceutically acceptable carrier or excipient therefor.

A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is

usually adapted for oral, parenteral, rectal administration or intravesical administration to the bladder and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusible solutions, suspensions or suppositories. Orally administrable compositions are generally preferred.

Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well known in normal pharmaceutical practice.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing.

Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.

5 The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. For systemic administration, dosage levels from 0.01mg to 100mg per kilogramme of body weight are useful in the treatment of pain. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 0.05 to 20, 20 to 250, or 0.1 to 500.0 mg, for  
10 example 0.2 to 5 and 0.1 to 250 mg; and such unit doses may be administered more than once a day, for example two or three a day, so that the total daily dosage is in the range of about 0.5 to 1000 mg; and such therapy may extend for a number of weeks or months.

15 No unacceptable toxicological effects are indicated with compounds of the invention when administered in accordance with the invention.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be  
20 incorporated by reference herein as though fully set forth.

The following Descriptions and Examples illustrate the preparation of the compounds of the invention.

## 25 **Abbreviations**

DMF = Dimethylformamide

BINAP = 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl

## **Description 1**

### 30 **5-Nitro-1-methylquinolinium iodide (D1)**

A mixture of 5-nitroquinoline (5 g, 0.028 mol) and iodomethane (5.4 ml, 0.086 mol) in DMF (8 ml) was heated to 40°C. After 2h a thick dark red precipitate was



formed. The reaction mixture was cooled and diluted with acetone. The solid was filtered and washed with acetone, and dried to give the title compound as an orange solid.

5 **Description 2**

5-Nitro-1-methyl-2-(1*H*)-quinolinone (D2)

5-Nitro-1-methylquinolinium iodide, (D1, 8.47 g, 0.027 mol) in warm water (90 ml) was added dropwise to a solution of potassium ferricyanide, (33.6 g, 0.1 mol) in 10% aqueous sodium hydroxide maintained at 45°C. After 5h the temperature was raised to 60°C and the solution heated for ca. 24h. The solution was cooled in an ice bath for 15 minutes and the grey-green solid filtered off, washed with water and dried. The crude solid was dissolved in a minimum amount of dichloromethane, filtered through silica gel and washed with ethyl acetate until no further product was eluted. Removal of ethyl acetate under reduced pressure afforded the title compound as a dark orange solid.

**Description 3**

5-Amino-1-methyl-2-(1*H*)-quinolinone (D3)

5-Nitro-1-methyl-2-(1*H*)-quinolinone, (D2, 2.59 g, 0.13 mol) in ethanol, (100 ml) and DMF (30 ml) was treated with 10% palladium on charcoal, (1 g, 50% w/w water). The reaction mixture was hydrogenated for a total of 8h at atmospheric pressure and then concentrated under reduced pressure. Trituration with ether afforded the title compound as a buff solid.

**Description 4**

1-(4-Chlorophenyl)piperidine-4-carboxylic acid, ethyl ester (D4)

Racemic BINAP (2.25 g, 0.0036 mol), palladium acetate (0.82 g, 3.65 mmol) and caesium carbonate (16.86 g, 0.051 mol) were suspended in 1,4-dioxane (100 ml) and sonicated for 45 minutes. 4-Bromo-chlorobenzene (5 g, 26.12 mmol) and

ethyl isonipecoacetate (4.11 g, 26.12 mmol) were added as a solution in 1,4-dioxane (100 ml). The mixture was heated to 105°C for 16h. On cooling the solvent was stripped off and the residues partitioned between water (100 ml) and diethyl ether (100 ml). The aqueous was re-extracted with ether. The combined  
5 layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash chromatography (ethylacetate/pet. ether eluent) yielded the title compound as a clear yellow oil.

### Description 5

10 1-(4-Chloro-phenyl)-piperidine-4-carboxylic acid (D5)

A solution of D4 (1.82 g, 0.0067 mol) in 1N LiOH (30 ml) and dioxane (30 ml) was stirred at room temperature for 16h and then evaporated *in vacuo*. Work up with 1M HCl and ethyl acetate gave the title compound as a yellow solid.

15

### Example 1

**1-(4-Chlorophenyl)-N-(1-methyl-2-oxo-1,2-dihydro-5-quinolinyl)-4-piperidinecarboxamide (E1)**



20 A suspension of 1-(4-chlorophenyl)-4-piperidinecarboxylic acid, (D5, 150mgs, 0.63 mM) in dichloromethane, (5 ml) under argon, was treated with oxalyl chloride, (0.164 ml, 1.88 mM) and 1 drop of DMF. After 2h, the solution was concentrated under high vacuum to remove excess reagents and then redissolved in dichloromethane, (10 ml). The solution was cooled in an ice-bath  
25 and treated with a solution of 5-amino-1-methyl-2-(1H)-quinolinone, D3 (109 mg, 0.03 mmol) and pyridine, (0.061 ml, 0.75 mmol). The reaction mixture was kept at room temperature for ca. 2h, then 45°C for a further 2h and 24h at room temperature again. The thick precipitate formed was removed by centrifugation, washed with dichloromethane then ether and dried to give the title compound as  
30 a buff solid, (156mgs, 63%). MH<sup>+</sup> 396.







**Example 2****4-(4-chlorophenyl)-*N*-5-quinolinylcyclohexanecarboxamide (E2)**








To a solution of 4-(4-chlorophenyl)cyclohexanecarboxylic acid (11.9 mg, 0.05 mmol) (Maybridge Chemical Company) in *N,N*-dimethylacetamide (1 mL) was added thionyl chloride (2.0 M solution in CH<sub>2</sub>Cl<sub>2</sub>, 25 μL, 0.05 mmol) and the resulting solution was stirred for 30 minutes before 5-aminoquinoline (7.2 mg, 0.05 mmol) (Aldrich Chemical Company) and diisopropylethylamine (19 μL, 0.15 mmol) were added as a solution in *N,N*-dimethylacetamide (0.5 mL). The solution was then stirred for 16 hours before the solvent was removed under reduced pressure and the residue purified by reverse phase preparative HPLC to yield a white solid (6.0 mg, 33%).  $MH^+ = 365$ .

Examples 3 - 40 presented in Table 1 were prepared by analogous procedures to those described in Examples 1 and 2.








**Table 1**









Example Number	Structure	$MH^+$ (observed)
3		379
4		407








5		406
6		385
7		396
8		380
9		381
10		394


11		394
12		360
13		428
14		394
15		371
16		448
17		414



18		380
19		394
20		414
21		398
22		414
23		414
24		382

25		398
26		430
27		416
28		364
29		360
30		376
31		425
32		435

33		477
34		431
35		401
36		408
37		366
38		384
39		408

40		363
----	--	-----

### Pharmacological Data

#### 5 (a) In vitro assay

As referenced above, the compounds of the invention are vanilloid receptor (VR1) antagonists and hence have useful pharmaceutical properties. Vanilloid receptor (VR1) antagonist activity can be confirmed and demonstrated for any particular compound by use of conventional methods, for example those  
 10 disclosed in standard reference texts such as D. Le Bars, M. Gozarin and S. W. Cadden, Pharmacological Reviews, 2001, 53(4), 597-652] or such other texts mentioned herein.

The screen used for the compounds of this invention was based upon a FLIPR based calcium assay, similar to that described by Smart et al. (British  
 15 Journal of Pharmacology, 2000, 129, 227-230).

Transfected astrocytoma 1321N1 cells, stably expressing human VR1, were seeded into FLIPR plates at 25,000cells/well (96-well plate) and cultured overnight.

20 The cells were subsequently loaded in medium containing 4 $\mu$ M Fluo-3 AM (Molecular Probes) for 2 hours, at room temperature, in the dark. The plates were then washed 4 times with Tyrode containing 1.5mM calcium, without probenecid.

25 The cells were pre-incubated with compound or buffer control at room temperature for 30 minutes. Capsaicin (Sigma) was then added to the cells. Compounds having antagonist activity against the human VR1 were identified by detecting differences in fluorescence when measured after capsaicin addition,


compared with no compound buffer controls. Thus, for example, in the buffer control capsaicin addition results in an increase in intracellular calcium concentration resulting in fluorescence. A compound having antagonist activity blocks the capsaicin binding to the receptor, there is no signalling and therefore no increase in intracellular calcium levels and consequently lower fluorescence. pKb values are generated from the IC<sub>50</sub> values using the Cheng-Prusoff equation.

All compounds tested by the above methodology had pKb > 6, preferred compounds having a pKb > 7.0.



Claims

1. A compound of formula (I),



(I)

or a pharmaceutically acceptable salt or solvate thereof, wherein,

- 10 P represents phenyl, quinoliny, isoquinoliny, 1,2,3,4-tetrahydroquinoliny, benzoisoxazolyl or benzothiazolyl;

P' represents phenyl, pyridiny or pyrimidiny;

R<sup>1</sup> and R<sup>3</sup> may be the same or different and represent alkyl, alkoxy, halo, -CF<sub>3</sub>, -OCF<sub>3</sub>, -OH, =O, -CN, -NO<sub>2</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -SO<sub>2</sub>R<sup>4</sup> or -NR<sup>4</sup>R<sup>5</sup>;

- 15 R<sup>2</sup> represents -H;

R<sup>4</sup> and R<sup>5</sup> may be the same or different and represent -H or alkyl;

m represents 0 or 1;


n represents 0, 1, 2, 3, 4 or 5; and

X represents N or CH.


- 20 2. A compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1, substantially as hereinbefore described with reference to any one of the Examples.

- 25 3. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1, which process comprises:

(a) reacting a compound of formula (II),



wherein, P, R<sup>1</sup> and n are as defined in relation to formula (I), with a compound of formula (III),



wherein, P', R<sup>3</sup>, m, n and X are as defined in relation to formula (I) and thereafter, as necessary, carrying out one or more of the following reactions:

- (i) converting one compound of formula (I) into another compound of formula (I);
- (ii) removing any protecting group;
- (iii) preparing a salt or a solvate of the compound so formed.


4. A pharmaceutical composition, which comprises a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1, and a pharmaceutically acceptable carrier or excipient therefor.

5. A compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1, for use as an active therapeutic substance.



*NR*

**PCT/EP2004/009078**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☒ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**